BEFORE USING OR HANDLING, READ
AND UNDERSTAND THIS LABEL

Indications:
For use in fish hatcheries for prevention and treatment of fungal infections on salmonid eggs.

Directions:
For prevention of fungal infections on salmonid eggs, apply PEROX-AID® at a concentration of 500 ppm for 15 minutes every other day.

For treatment of existing fungal infections on salmonid eggs, apply PEROX-AID® at a concentration of 500 ppm for 60 minutes every other day.

Dosage:
Prior to use on salmon eggs dilute PEROX-AID® to 500 ppm based on 100% active ingredient.

Calculations for the amount of PEROX-AID® to be added to the water inflow into the egg holding apparatus is dependent on the rate of water flow. A water medicator may be used to perform this function. Use 1.43 mL of PEROX-AID® per litre of water treated.

Storage:
- Avoid outside storage
- Store in original container in dry location
- Keep container out of sun and away from heat

Syndel
CANADA

MANUFACTURED FOR/
FABRIQUÉ POUR:
SYNDEL CANADA
NANAIMO, BC
CANADA V9S 4M9
ver. 122616

For Chemical Emergency Spill, Leak, Fire, Exposure or Accident
CALL DAY OR NIGHT:
US - CHEMTREC 1-800-424-9300
CANADA - CANUTEC (collect) 1-613-996-6666

UN2014,
HYDROGEN PEROXIDE 35%, Aqueous Solutions,
5.1, (8), PGII, ERG# 140

PEROX-AID®

HYDROGEN PEROXIDE 35% W/W, AQUEOUS SOLUTION
PEROXYDE D' HYDROGÈNE 35% P/P, SOLUTION AQUEUSE

Veterinary Use Only
Usage vétérinaire seulement

ANTIFUNGAL FOR USE ON SALMONID EGGS ONLY
(ANTIFONGIQUE POUR USAGE SUR OEUFS DE SAUMONS SEULEMENT)

DIN 02238749

Read Entire Label Before Using This Product
Net Contents 208 Litres (55 (U.S.) GALLONS)
Bien lire les instructions avant d'utiliser ce produit
Contenu net 208 Litres (55 GALLONS U.S.)

ACTIVE INGREDIENT/INGRÉDIENT ACTIF
Hydrogen peroxide (H_2O_2) I Peroxyde d' hydrogène (H_2O_2) 35% w/w

WARNINGS!
HAZARDOUS INGREDIENTS

1. Strong oxidizer - Handle with care to avoid spillage
2. Do not expose to heat or open flame
3. Wear safety glasses and impervious gloves when handling this product
4. End users are required to comply with all safety and enviromnental regulations
5. Keep out of reach of children

MISES EN GARDE
INGRÉDIENTS NOCIFS

1. Agent oxidant puissant - Manipuler avec soins, éviter les renversements.
2. Ne pas exposer à la chaleurs ou aux flammes nues.
3. Toujours porter des lunettes et gants de sécurité pour manipuler ce produit.
4. Toujours respecter les normes de sécurité et environnementales appropriées lorsque vous utilisez ce produit.
5. Garder hors de la portée des enfants.

ASSUREZ VOUS DE BIEN LIRE ET COMPRENDRE LES
INSTRUCTIONS CI-APRÈS AVANT D'UTILISER CE PRODUIT.

Indications:
Pour utilisation dans l'industrie de la pisciculture en tant qu'agent fongicide comme moyen de prévention ou de traitement pour les oeufs de saumons.

Mode d'emploi:
Pour prévenir d'infection sur des oeufs de saumons, appliquer PEROX-AID® à une concentration de 500 ppm pour 15 minutes à tous les deux jours.

Pour traitement d'infections existantes pour oeufs de saumons, appliquer PEROX-AID® à une concentration de 500 ppm pour 60 minutes à tous les deux jours.

Posologie:
Avant d'utiliser PEROX-AID® sur des oeufs de saumons, diluer le produit avec de l'eau à une concentration de 500 ppm sur une base d'agent actif pur.

Le calcul de la quantité de PEROX-AID® qui doit être ajoutée à l'eau entrant dans le réservoir qui contient les oeufs, dépend du débit. Un appareil de dosage pour le traitement d'eau ("water medicator") peut être utilisé pour accomplir cette tâche. Utiliser 1,43 mL de PEROX-AID® par litre d'eau à traiter.

Entreposage:
- Éviter d'entreposer le produit à l'extérieur
- Entreposer le produit dans le contenant original dans un endroit sec
- Garder le contenant loin soleil et de la chaleur

LOT NO:

EXPIRATION DATE I DATE D'EXPIRATION:

Information Only